

Some Tree-Star Ramsey Numbers

E. J. COCKAYNE

University of Victoria, Victoria, British Columbia, Canada

Communicated by Frank Harary

Received November 13, 1973

The Ramsey Number $r(G_1, G_2)$ is the least integer N such that for every graph G with N vertices, either G has the graph G_1 as a subgraph or \bar{G} , the complement of G , has the graph G_2 as a subgraph.

In this paper we embed the paths P_m in a much larger class \mathcal{T} of trees and then show how some evaluations by T. D. Parsons of Ramsey numbers $r(P_m, K_{1,n})$, where $K_{1,n}$ is the star of degree n , are also valid for $r(T_m, K_{1,n})$ where $T_m \in \mathcal{T}$.

1. INTRODUCTION

The quantity $r(G_1, G_2)$ is the least integer N such that for every graph G with N vertices, either G has the graph G_1 as a subgraph or \bar{G} , the complement of G , has the graph G_2 as a subgraph.

The well-known theorem of F. Ramsey [4] ensures the existence of the numbers $r(G_1, G_2)$, and many authors are currently evaluating these "Ramsey numbers" for special G_1, G_2 . The reader is referred to the excellent survey article by S. Burr [1] for an extensive bibliography.

T. D. Parsons [3] has recently evaluated $r(P_m, K_{1,n})$ where P_m is a path with m vertices and $K_{1,n}$ is the star of degree n . In Section 2 of this note we embed the paths in a large class \mathcal{T} of trees and then in Section 3 show that many of Parsons' results are also true if P_m is replaced by T_m where T_m is any m -vertex member of \mathcal{T} . Most of the proofs in Section 3 are identical to those of Parsons and therefore, we do not, repeat them.

Any undefined term or notation is given in Harary's book [2].

2. THE CLASS OF TREES \mathcal{T}

LEMMA 1. *If a graph G has $\delta(G) \geq k$, then G contains as a subgraph any tree on $k + 1$ vertices.*

Proof. By induction on k . The result is true for $k = 1$ and 2. Now consider any graph G with $\delta(G) \geq k + 1$ and any tree T having $k + 2$ vertices. Delete from T an edge $[v, w]$ and vertex w which has degree 1

in T , thus forming a tree T' with $k + 1$ vertices. Delete from G any vertex and all edges adjacent to this vertex, forming G' with $\delta(G') \geq k$. By the induction hypothesis, $G' \supset T'$. If $d(x, F)$ denotes the degree of vertex x in the graph F , then $d(v, T') \leq k$ and $d(v, G) \geq k + 1$. Hence v is adjacent to a vertex of G not in T' and the addition of such an edge produces $T \subset G$, thus completing the proof.

Let T be a spanning tree of G which has m vertices. A permutation π of $V(G)$ is T -preserving if and only if for each $[\alpha, \beta] \in E(T)$, $[\pi(\alpha), \pi(\beta)] \in E(G)$. Let x be a vertex of G and X be the set of all images of x under T -preserving permutations of $V(G)$.

We say that (T, x) is a *complete pair* if and only if the assumption that each $x \in X$ has degree $m - 1$ in G implies that G is the complete graph K_m .

As an illustration, consider the pair $(T, 1)$ of Fig. 1:




FIGURE 1.

Since $1 \in X$, $d(1, G) = 4$. Therefore

$1 \rightarrow 3$	$1 \rightarrow 2$	$1 \rightarrow 4$
$2 \rightarrow 2$	$2 \rightarrow 3$	$2 \rightarrow 2$
$3 \rightarrow 5$	$3 \rightarrow 5$	$3 \rightarrow 5$
$4 \rightarrow 1$	$4 \rightarrow 1$	$4 \rightarrow 1$
$5 \rightarrow 4$	$5 \rightarrow 4$	$5 \rightarrow 3$

are T -preserving permutations. Hence the vertices 2, 3, 4 are in X and have degree 4 in G . Hence $G = K_5$ and $(T, 1)$ is a complete pair.

The trees P_{2n+1} and $K_{1,n}$ together with their respective center vertices do not form complete pairs. Further, the tree with vertex set

$$\{x_1\} \cup \{x_2\} \cup Y_1 \cup Y_2, \quad \text{where } |Y_1|, |Y_2| \geq 2,$$

and edge set $\{[x_1, x_2]\} \cup \{[x_1, y]: y \in Y_1\} \cup \{[x_2, y]: y \in Y_2\}$ does not form a complete pair with either of the vertices x_1, x_2 . We have not as yet completely characterized complete pairs. The following result shows that degree-one vertices always yield complete pairs.

LEMMA 2. *If x has degree one in a spanning tree T of G , then (T, x) is a complete pair.*

Proof. Suppose $|V(G)| = m$. Let the length of the path from vertex v to x be $D(v)$. It is sufficient to prove that for all k , each vertex v with $D(v) = k$ is an image of x under a T -preserving permutation. We do this by induction on k . The case $k = 0$ is trivial. Now assume the result for all $k \leq j$ and suppose that $D(w) = j + 1$. Let the vertices in the path from x to w be x, x_1, \dots, x_j, w . By the induction hypothesis, x, x_1, \dots, x_j have degree $m - 1$ in G . Therefore the permutation which maps the ordered set $\{x, x_1, \dots, x_j, w\}$ into $\{w, x, x_1, \dots, x_j\}$ and leaves all other vertices invariant is T -preserving and the image of x is w as required. This completes the proof.

A tree T is in the class \mathcal{T} if and only if some vertex v of degree one in T and its incident edge $[v, w]$ may be deleted forming a tree T' where (T', w) is a complete pair.

Lemma 2 implies that \mathcal{T} contains all trees which have a vertex of degree one adjacent to vertex of degree two, hence in particular \mathcal{T} contains all paths P_m . However, there are many trees in \mathcal{T} without this property. No star $K_{1,n}$ for $n \geq 3$ is in \mathcal{T} . For $m \leq 7$, the only tree not in \mathcal{T} which is not a star is that shown in Fig. 2.




FIG. 2. A tree not in \mathcal{T} .

A quick glance at the tree diagrams of [2, p. 233] shows there are at most five trees not in \mathcal{T} among the 23 8-vertex trees.

3. RAMSEY NUMBERS

The following two results have recently appeared in the literature. Let T_m be any tree with m vertices.

THEOREM 1 (Burr [1]).

$$r(T_m, K_{1,n}) \leq m + n - 1.$$

THEOREM 2 (Burr [1]). *If $m - 1 \mid n - 1$, then*

$$r(T_m, K_{1,n}) = m + n - 1.$$

The next result generalizes Lemma 4 in Parsons' work [3].

LEMMA 3. *If $T_m \in \mathcal{T}$ and $n \not\equiv 1 \pmod{m-1}$, then*

$$r(T_m, K_{1,n}) \leq m + n - 2.$$

Proof. By induction on n . The statement is true for $n = 1$. Suppose $n > 1$, $n \not\equiv 1 \pmod{m-1}$, and G has $m + n - 2$ vertices. Suppose $G \not\supset T_m$ and $\Delta(\bar{G}) \leq n - 1$. Then

$$\delta(G) = m + n - 3 - \Delta(\bar{G}) \geq m + n - 3 - (n - 1) = m - 2,$$

and so by definition of \mathcal{T} and Lemma 1, G contains a tree T'_{m-1} formed from T_m by removal of a vertex v of degree one in T_m and its incident edge $[v, w]$, and such that (T'_{m-1}, w) is a complete pair. Let the vertex set of T'_{m-1} be $W = \{v_1, \dots, v_{m-1}\}$ and the remaining vertices be $H = \{v_m, \dots, v_{m+n-2}\}$. If the image of w under any T'_{m-1} -preserving permutation of W is adjacent in G to any vertex of H , $G \supset T_m$. Hence there is no such adjacency, and each such image, having degree $\geq m - 2$ in G , has degree $m - 2$ in $\langle W \rangle$, the subgraph induced by W in G . Since (T'_{m-1}, w) is a complete pair, $G = K_{m-1} \cup \langle H \rangle$. If $n \leq m - 1$, then $d(v_m, \bar{G}) \geq m - 1 \geq n$, hence $n > m - 1$. By the induction hypothesis, since $n - m + 1 \not\equiv 1 \pmod{m-1}$, $r(T_m, K_{1,n-m+1}) \leq m + (n - m + 1) - 2 = n - 1$. But H has $n - 1$ vertices and $H \not\supset T_m$ so $\Delta(\bar{H}) \geq n - m + 1$. Then

$$\Delta(\bar{G}) \geq \Delta(\bar{H}) + (m - 1) \geq n - m + 1 + m - 1 = n,$$

a contradiction. The lemma follows.

LEMMA 4. *If F_m is any connected graph with m vertices and $n \equiv 0, 2 \pmod{m-1}$, then*

$$r(F_m, K_{1,n}) > m + n - 3.$$

Proof. Identical to that of Theorem 4 in [3].

THEOREM 3. *If $T_m \in \mathcal{T}$ and $m \equiv 0, 2 \pmod{m-1}$, then*

$$r(T_m, K_{1,n}) = m + n - 2.$$

Proof. Immediate from Lemmas 3 and 4.

THEOREM 4. If $T_m \in \mathcal{T}$, $n \not\equiv 1 \pmod{m-1}$, and $n \geq (m-3)^2$, then

$$r(T_m, K_{1,n}) = m + n - 2.$$

Proof. The same arguments used in the proofs of Lemma 5 and Theorem 5 in [3] establish that for any connected graph F_m with m vertices, where m satisfies the hypotheses, $r(F_m, K_{1,n}) > m + n - 3$, and the result follows from Lemma 3.

THEOREM 5. If $T_m \in \mathcal{T}$, $n \not\equiv 1 \pmod{m-1}$, and $n \equiv 1 \pmod{m-2}$, then

$$r(T_m, K_{1,n}) = m + n - 2.$$

Proof. Let T'_{m-1} be any tree formed from T_m by deleting a vertex of degree one. Then using Theorem 2,

$$r(T_m, K_{1,n}) \geq r(T'_{m-1}, K_{1,n}) = (m-1) + n - 1.$$

The result now follows from Lemma 3.

THEOREM 6. If $T_m \in \mathcal{T}$, $m > 3$, $n \equiv -1 \pmod{m-1}$, and $n \geq m-1$, then

$$r(T_m, K_{1,n}) = m + n - 2.$$

Proof. Identical to that of Theorem 7 in [3].

ACKNOWLEDGMENT

The author gratefully acknowledges the support of the Canadian National Research Council Grant No. A7544.

REFERENCES

1. S. BURR, Generalized Ramsey theory for graphs—A survey, to appear in "Graphs and Combinatorics" (R. Bari and F. Harary, Eds.), Springer-Verlag, Berlin, 1974.
2. F. HARARY, "Graph Theory," Addison-Wesley, Reading, Mass., 1969.
3. T. D. PARSONS, Path-star Ramsey numbers, *Proc. Amer. Math. Soc.* to appear.
4. F. RAMSEY, On a problem of formal logic, *Proc. London Math. Soc.* **30** (1930), 264–286.